Chapter 05.10 Shortest Path of a Robot

After reading this chapter, you should be able to:

- 1. find the shortest smooth path through consecutive points, and
- 2. compare the lengths of different paths.

Example

Peter: "Dr. Kaw, I am taking a course in manufacturing. We are solving the following problem. A robot arm with a rapid laser is used to do a quick quality check, such as the radius of hole, on six holes on a rectangular plate $15"\times10"$ at several points as shown in Table 1 and Figure 1.

Table 1 The coordinate values of six holes on a rectangle plate.

x	У
2.00	7.2
4.5	7.1
5.25	6.0
7.81	5.0
9.20	3.5
10.60	5.0

I am using Excel to fit a fifth order polynomial through the 6 points. But, when I plot the polynomial, it is taking a long path! (Figure 2)"

Kaw: "Why do you not just join the consecutive points by a straight line; just like the kids do at Pizza HutTM with those 'Connect the dots' activities?"

Peter: "You are making me hungry and I wish it were that easy. The path of the robot going from one point to another point needs to be smooth so as to avoid sharp jerks in the arm that can otherwise create premature wear and tear of the robot arm."

Kaw: "As I recall, you took my course in Numerical Methods. What was that one year ago?"

Peter: "Yes, your memory is sharp, but my retention from that course – can we not talk about that?!?"

Kaw: "Come into my office. I wrote this program using Maple. See this function, $f(x) = 1/(1+25x^2)$. I am choosing 7 points equidistantly (Table 2) between -1 and 1.

Figure 1 Locations of holes on the rectangular plate.



Figure 2 Approximating the path of the robot using 5th order polynomial.

Now look at the sixth order interpolating polynomial and the original function (Figure 3). See the oscillations in the interpolating polynomial. In 1901, Runge used this example function to show that higher order interpolation is a bad idea. One of the solutions to your

robot path problem is to use quadratic or cubic spline interpolation. That will give you a smooth curve with fewer oscillations, and a smoother and shorter path."

0.038462

0.264706

0.264706

0.082569 0.0385

0.0826

1

Table 2	The coordinate values	of 7 equidistantly spaced points				
		x	$y = \frac{1}{1 + 25x^2}$			

-0.66667

-0.33333

0.333333

0.666667

-1

0

1



Figure 3 Runge's function interpolated.

Peter: "Okay. Let's give that a try." **Kaw:** "Now, let's try generating a set of cubic splines to go through the data:"



Figure 4 Path of the robot arm using cubic spline interpolation.

Peter: "Wow! That (Figure 4) looks much better!"

Kaw: "It may look better, but let's find out for sure. See if you can combine the two plots (Figure 5) and compare the lengths of each path."

Peter: "The length of a path *S* if y = f(x) from *a* to *b* is given by

$$S = \int_{a}^{b} \sqrt{1 + \left(\frac{df}{dx}\right)^2} \, dx$$

Right?"

Table 3Comparison of the length of curves.

<u></u>	•••= ••••	
Type of interpolation	5th order polynomial	Cubic Spline
Length of Curve	14.919"	11.248"

Kaw: "Yes! You solved the problem. See Table 3 for answers."

Peter: "I guess your class was good for something after all, Dr. Kaw."

Kaw: "Are you sure? You could have always fallen back on the connecting-the-dots method. Besides, you don't want to grow up ... you're a Pizza Hut[™] kid, right?" **Peter:** "That's a Toys Я' Us[™] kid. You'll do anything to be reminded of songs, won't you?"



Figure	5	Path	of	robot	arm	compared	using	polynomial	interpolation	and	cubic	spline
interpol	atio	on.										

INTERPOLATION				
Topic	Shortest path of a robot			
Summary	An example of interpolation: A robot arm path needs to be developed			
	over several points on a flat plate. The path needs to be smooth to avoid			
	sudden jerky motion and at the same time needs to be short.			
Major	General Engineering			
Authors	Autar Kaw, Michael Keteltas			
Date	December 7, 2008			
Web Site	http://numericalmethods.eng.usf.edu			