

Gaussian Elimination

Industrial Engineering Majors

Author(s): Autar Kaw

<http://numericalmethods.eng.usf.edu>

Transforming Numerical Methods Education for STEM
Undergraduates

Naïve Gauss Elimination

<http://numericalmethods.eng.usf.edu>

Naïve Gaussian Elimination

A method to solve simultaneous linear equations of the form $[A][X]=[C]$

Two steps

1. Forward Elimination
2. Back Substitution

Forward Elimination

The goal of forward elimination is to transform the coefficient matrix into an upper triangular matrix

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$



$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$

Forward Elimination

A set of n equations and n unknowns

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n = b_2$$

⋮
⋮
⋮
⋮

$$a_{n1}x_1 + a_{n2}x_2 + a_{n3}x_3 + \dots + a_{nn}x_n = b_n$$

(n-1) steps of forward elimination

Forward Elimination

Step 1

For Equation 2, divide Equation 1 by a_{11} and multiply by a_{21} .

$$\left[\frac{a_{21}}{a_{11}} \right] (a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1)$$

$$a_{21}x_1 + \frac{a_{21}}{a_{11}}a_{12}x_2 + \dots + \frac{a_{21}}{a_{11}}a_{1n}x_n = \frac{a_{21}}{a_{11}}b_1$$

Forward Elimination

Subtract the result from Equation 2.

$$\begin{array}{rcl} a_{21}x_1 + a_{22}x_2 + a_{23}x_3 + \dots + a_{2n}x_n & = & b_2 \\ - \quad a_{21}x_1 + \frac{a_{21}}{a_{11}}a_{12}x_2 + \dots + \frac{a_{21}}{a_{11}}a_{1n}x_n & = & \frac{a_{21}}{a_{11}}b_1 \\ \hline \left(a_{22} - \frac{a_{21}}{a_{11}}a_{12} \right)x_2 + \dots + \left(a_{2n} - \frac{a_{21}}{a_{11}}a_{1n} \right)x_n & = & b_2 - \frac{a_{21}}{a_{11}}b_1 \end{array}$$

or $\overset{'}{a_{22}x_2} + \dots + \overset{'}{a_{2n}x_n} = \overset{'}{b_2}$

Forward Elimination

Repeat this procedure for the remaining equations to reduce the set of equations as

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\dot{a_{32}}x_2 + \dot{a_{33}}x_3 + \dots + \dot{a_{3n}}x_n = \dot{b_3}$$

$$\begin{matrix} \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot \end{matrix}$$

$$\dot{a_{n2}}x_2 + \dot{a_{n3}}x_3 + \dots + \dot{a_{nn}}x_n = \dot{b_n}$$

End of Step 1

Forward Elimination

Step 2

Repeat the same procedure for the 3rd term of Equation 3.

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$^{''}a_{33}x_3 + \dots + ^{''}a_{3n}x_n = ^{''}b_3$$

$$\begin{matrix} \cdot & \cdot \\ \cdot & \cdot \\ \cdot & \cdot \end{matrix}$$

$$^{'''}a_{n3}x_3 + \dots + ^{'''}a_{nn}x_n = ^{'''}b_n$$

End of Step 2

Forward Elimination

At the end of (n-1) Forward Elimination steps, the system of equations will look like

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_{3n}}x_n = \ddot{b_3}$$

.

.

.

$$a_{nn}^{(n-1)}x_n = b_n^{(n-1)}$$

End of Step (n-1)

Matrix Form at End of Forward Elimination

$$\begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a'_{22} & a'_{23} & \cdots & a'_{2n} \\ 0 & 0 & a''_{33} & \cdots & a''_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & a_{nn}^{(n-1)} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b'_2 \\ b''_3 \\ \vdots \\ b_n^{(n-1)} \end{bmatrix}$$

Back Substitution

Solve each equation starting from the last equation

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ -96.21 \\ 0.735 \end{bmatrix}$$

Example of a system of 3 equations

Back Substitution Starting Eqns

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_n}x_n = \ddot{b_3}$$

. . .
.

$$a_{nn}^{(n-1)}x_n = b_n^{(n-1)}$$

Back Substitution

Start with the last equation because it has only one unknown

$$x_n = \frac{b_n^{(n-1)}}{a_{nn}^{(n-1)}}$$

Back Substitution

$$x_n = \frac{b_n^{(n-1)}}{a_{nn}^{(n-1)}}$$

$$x_i = \frac{b_i^{(i-1)} - a_{i,i+1}^{(i-1)}x_{i+1} - a_{i,i+2}^{(i-1)}x_{i+2} - \dots - a_{i,n}^{(i-1)}x_n}{a_{ii}^{(i-1)}} \text{ for } i = n-1, \dots, 1$$

$$x_i = \frac{b_i^{(i-1)} - \sum_{j=i+1}^n a_{ij}^{(i-1)}x_j}{a_{ii}^{(i-1)}} \text{ for } i = n-1, \dots, 1$$

THE END

<http://numericalmethods.eng.usf.edu>

Naïve Gauss Elimination Example

<http://numericalmethods.eng.usf.edu>

Example: Production Optimization

To find the number of toys a company should manufacture per day to optimally use their injection-molding machine and the assembly line, one needs to solve the following set of equations. The unknowns are the number of toys for boys, x_1 , number of toys for girls, x_2 , and the number of unisexual toys, x_3 .

$$\begin{bmatrix} 0.3333 & 0.1667 & 0.6667 \\ 0.1667 & 0.6667 & 0.3333 \\ 1.05 & -1.00 & 0.00 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 756 \\ 1260 \\ 0 \end{bmatrix}$$

Find the values of x_1 , x_2 , and x_3 using Naïve Gauss Elimination.

Example: Production Optimization

Forward Elimination: Step 1

$$Row2 - \left[\frac{Row1}{0.3333} \right] \times (0.1667) =$$

Yields

$$\begin{bmatrix} 0.3333 & 0.1667 & 0.6667 \\ 0 & 0.58332 & -0.00015002 \\ 1.05 & -1.00 & 0.00 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 756 \\ 881.89 \\ 0 \end{bmatrix}$$

Example: Production Optimization

Forward Elimination: Step 1

$$Row3 - \left[\frac{Row1}{0.3333} \right] \times (1.05) =$$

Yields

$$\begin{bmatrix} 0.3333 & 0.1667 & 0.6667 \\ 0 & 0.58332 & -0.00015002 \\ 0 & -1.5252 & -2.1003 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 756 \\ 881.89 \\ -2381.6 \end{bmatrix}$$

Example: Production Optimization

Forward Elimination: Step 2

$$Row3 - \left[\frac{Row2}{0.58332} \right] \times (-1.5252) =$$

Yields

$$\begin{bmatrix} 0.3333 & 0.1667 & 0.6667 \\ 0 & 0.58332 & -0.00015002 \\ 0 & 0 & -2.1007 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 756 \\ 881.89 \\ -75.864 \end{bmatrix}$$

This is now ready for Back Substitution

Example: Production Optimization

Back Substitution: Solve for x_3 using the third equation

$$-2.1007x_3 = -75.864$$

$$\begin{aligned}x_3 &= \frac{-75.864}{-2.1007} \\&= 36.113\end{aligned}$$

Example: Production Optimization

Back Substitution: Solve for x_2 using the second equation

$$0.58332x_2 + (-0.00015002)x_3 = 881.89$$

$$\begin{aligned}x_2 &= \frac{881.89 - (-0.00015002)x_3}{0.58332} \\&= \frac{881.89 - (-0.00015002) \times 36.113}{0.58332} \\&= 1511.8\end{aligned}$$

Example: Production Optimization

Back Substitution: Solve for x_1 using the first equation

$$0.3333x_1 + 0.1667x_2 + 0.6667x_3 = 756$$

$$\begin{aligned}x_1 &= \frac{756 - 0.1667x_2 - 0.6667x_3}{0.3333} \\&= \frac{756 - 0.1667 \times 1511.8 - 0.6667 \times 36.113}{0.3333} \\&= 1439.8\end{aligned}$$

Example: Production Optimization

Solution:

The solution vector is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1439.8 \\ 1511.8 \\ 36.113 \end{bmatrix}$$

1440 toys for boys should be produced

1512 toys for girls should be produced

36 unisexual toys should be produced

THE END

<http://numericalmethods.eng.usf.edu>

Naïve Gauss Elimination Pitfalls

<http://numericalmethods.eng.usf.edu>

Pitfall#1. Division by zero

$$10x_2 - 7x_3 = 3$$

$$6x_1 + 2x_2 + 3x_3 = 11$$

$$5x_1 - x_2 + 5x_3 = 9$$

$$\begin{bmatrix} 0 & 10 & -7 \\ 6 & 2 & 3 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 3 \\ 11 \\ 9 \end{bmatrix}$$

Is division by zero an issue here?

$$12x_1 + 10x_2 - 7x_3 = 15$$

$$6x_1 + 5x_2 + 3x_3 = 14$$

$$5x_1 - x_2 + 5x_3 = 9$$

$$\begin{bmatrix} 12 & 10 & -7 \\ 6 & 5 & 3 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 14 \\ 9 \end{bmatrix}$$

Is division by zero an issue here?

YES

$$12x_1 + 10x_2 - 7x_3 = 15$$

$$6x_1 + 5x_2 + 3x_3 = 14$$

$$24x_1 - x_2 + 5x_3 = 28$$

$$\begin{bmatrix} 12 & 10 & -7 \\ 6 & 5 & 3 \\ 24 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 14 \\ 28 \end{bmatrix} \rightarrow \begin{bmatrix} 12 & 10 & -7 \\ 0 & 0 & 6.5 \\ 12 & -21 & 19 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 15 \\ 6.5 \\ -2 \end{bmatrix}$$

Division by zero is a possibility at any step
of forward elimination

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Exact Solution

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Solve it on a computer using 6 significant digits with chopping

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.9625 \\ 1.05 \\ 0.999995 \end{bmatrix}$$

Pitfall#2. Large Round-off Errors

$$\begin{bmatrix} 20 & 15 & 10 \\ -3 & -2.249 & 7 \\ 5 & 1 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 45 \\ 1.751 \\ 9 \end{bmatrix}$$

Solve it on a computer using 5 significant digits with chopping

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0.625 \\ 1.5 \\ 0.99995 \end{bmatrix}$$

Is there a way to reduce the round off error?

Avoiding Pitfalls

Increase the number of significant digits

- Decreases round-off error
- Does not avoid division by zero

Avoiding Pitfalls

Gaussian Elimination with Partial Pivoting

- Avoids division by zero
- Reduces round off error

THE END

<http://numericalmethods.eng.usf.edu>

Gauss Elimination with Partial Pivoting

<http://numericalmethods.eng.usf.edu>

Pitfalls of Naïve Gauss Elimination

- Possible division by zero
- Large round-off errors

Avoiding Pitfalls

Increase the number of significant digits

- Decreases round-off error
- Does not avoid division by zero

Avoiding Pitfalls

Gaussian Elimination with Partial Pivoting

- Avoids division by zero
- Reduces round off error

What is Different About Partial Pivoting?

At the beginning of the k^{th} step of forward elimination,
find the maximum of

$$|a_{kk}|, |a_{k+1,k}|, \dots, |a_{nk}|$$

If the maximum of the values is $|a_{pk}|$
in the p^{th} row, $k \leq p \leq n$, then switch rows p and k .

Matrix Form at Beginning of 2nd Step of Forward Elimination

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & a_{n2} & a_{n3} & a_{n4} & a_{nn} \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{array} \right] = \left[\begin{array}{c} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{array} \right]$$

Example (2nd step of FE)

$$\begin{bmatrix} 6 & 14 & 5.1 & 3.7 & 6 \\ 0 & -7 & 6 & 1 & 2 \\ 0 & 4 & 12 & 1 & 11 \\ 0 & 9 & 23 & 6 & 8 \\ 0 & -17 & 12 & 11 & 43 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} 5 \\ -6 \\ 8 \\ 9 \\ 3 \end{bmatrix}$$

Which two rows would you switch?

Example (2nd step of FE)

$$\left[\begin{array}{ccccc} 6 & 14 & 5.1 & 3.7 & 6 \\ 0 & -17 & 12 & 11 & 43 \\ 0 & 4 & 12 & 1 & 11 \\ 0 & 9 & 23 & 6 & 8 \\ 0 & -7 & 6 & 1 & 2 \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{array} \right] = \left[\begin{array}{c} 5 \\ 3 \\ 8 \\ 9 \\ -6 \end{array} \right]$$

Switched Rows

Gaussian Elimination with Partial Pivoting

A method to solve simultaneous linear equations of the form $[A][X]=[C]$

Two steps

1. Forward Elimination
2. Back Substitution

Forward Elimination

Same as naïve Gauss elimination method except that we switch rows before **each** of the $(n-1)$ steps of forward elimination.

Example: Matrix Form at Beginning of 2nd Step of Forward Elimination

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & a_{n2} & a_{n3} & a_{n4} & a_{nn} \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{array} \right] = \left[\begin{array}{c} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{array} \right]$$

Matrix Form at End of Forward Elimination

$$\left[\begin{array}{cccc|c} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ 0 & a_{22} & a_{23} & \cdots & a_{2n} \\ 0 & 0 & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \cdots & \vdots \\ 0 & 0 & 0 & 0 & a_{nn}^{(n-1)} \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \\ \vdots \\ x_n \end{array} \right] = \left[\begin{array}{c} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n^{(n-1)} \end{array} \right]$$

Back Substitution Starting Eqns

$$a_{11}x_1 + a_{12}x_2 + a_{13}x_3 + \dots + a_{1n}x_n = b_1$$

$$\dot{a_{22}}x_2 + \dot{a_{23}}x_3 + \dots + \dot{a_{2n}}x_n = \dot{b_2}$$

$$\ddot{a_{33}}x_3 + \dots + \ddot{a_n}x_n = \ddot{b_3}$$

. . .
.

$$a_{nn}^{(n-1)}x_n = b_n^{(n-1)}$$

Back Substitution

$$x_n = \frac{b_n^{(n-1)}}{a_{nn}^{(n-1)}}$$

$$x_i = \frac{b_i^{(i-1)} - \sum_{j=i+1}^n a_{ij}^{(i-1)} x_j}{a_{ii}^{(i-1)}} \text{ for } i = n-1, \dots, 1$$

THE END

<http://numericalmethods.eng.usf.edu>

Gauss Elimination with Partial Pivoting Example

<http://numericalmethods.eng.usf.edu>

Example 2

Solve the following set of equations
by Gaussian elimination with partial
pivoting

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

Example 2 Cont.

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix} \Rightarrow \begin{bmatrix} 25 & 5 & 1 & : & 106.8 \\ 64 & 8 & 1 & : & 177.2 \\ 144 & 12 & 1 & : & 279.2 \end{bmatrix}$$

1. Forward Elimination
2. Back Substitution

Forward Elimination

Number of Steps of Forward Elimination

Number of steps of forward elimination is
 $(n-1) = (3-1) = 2$

Forward Elimination: Step 1

- Examine absolute values of first column, first row and below.

$|25|, |64|, |144|$

- Largest absolute value is 144 and exists in row 3.
- Switch row 1 and row 3.

$$\left[\begin{array}{ccc|c} 25 & 5 & 1 & : 106.8 \\ 64 & 8 & 1 & : 177.2 \\ 144 & 12 & 1 & : 279.2 \end{array} \right] \Rightarrow \left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 64 & 8 & 1 & : 177.2 \\ 25 & 5 & 1 & : 106.8 \end{array} \right]$$

Forward Elimination: Step 1 (cont.)

$$\begin{bmatrix} 144 & 12 & 1 & : & 279.2 \\ 64 & 8 & 1 & : & 177.2 \\ 25 & 5 & 1 & : & 106.8 \end{bmatrix}$$

Divide Equation 1 by 144 and multiply it by 64, $\frac{64}{144} = 0.4444$.

$$[144 \ 12 \ 1 \ : \ 279.2] \times 0.4444 = [63.99 \ 5.333 \ 0.4444 \ : \ 124.1]$$

Subtract the result from
Equation 2

$$\begin{array}{r} [64 \qquad \qquad \qquad 1 \ : \ 177.2] \\ - [63.99 \qquad 5.333 \qquad 0.4444 \ : \ 124.1] \\ \hline [0 \qquad 2.667 \qquad 0.5556 \ : \ 53.10] \end{array}$$

Substitute new equation for
Equation 2

$$\begin{bmatrix} 144 & 12 & 1 & : & 279.2 \\ 0 & 2.667 & 0.5556 & : & 53.10 \\ 25 & 5 & 1 & : & 106.8 \end{bmatrix}$$

Forward Elimination: Step 1 (cont.)

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.667 & 0.5556 & : 53.10 \\ 25 & 5 & 1 & : 106.8 \end{array} \right] \quad \text{Divide Equation 1 by 144 and multiply it by 25, } \frac{25}{144} = 0.1736.$$

$$[144 \ 12 \ 1 \ : \ 279.2] \times 0.1736 = [25.00 \ 2.083 \ 0.1736 \ : \ 48.47]$$

Subtract the result from
Equation 3

$$\begin{array}{r} [25 \ 5 \ 1 \ : \ 106.8] \\ - [25 \ 2.083 \ 0.1736 \ : \ 48.47] \\ \hline [0 \ 2.917 \ 0.8264 \ : \ 58.33] \end{array}$$

Substitute new equation for
Equation 3

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.667 & 0.5556 & : 53.10 \\ 0 & 2.917 & 0.8264 & : 58.33 \end{array} \right]$$

Forward Elimination: Step 2

- Examine absolute values of second column, second row and below.

$$|2.667|, |2.917|$$

- Largest absolute value is 2.917 and exists in row 3.
- Switch row 2 and row 3.

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.667 & 0.5556 & : 53.10 \\ 0 & 2.917 & 0.8264 & : 58.33 \end{array} \right] \Rightarrow \left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.917 & 0.8264 & : 58.33 \\ 0 & 2.667 & 0.5556 & : 53.10 \end{array} \right]$$

Forward Elimination: Step 2 (cont.)

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.917 & 0.8264 & : 58.33 \\ 0 & 2.667 & 0.5556 & : 53.10 \end{array} \right]$$

Divide Equation 2 by 2.917 and multiply it by 2.667,
 $\frac{2.667}{2.917} = 0.9143$.

$$[0 \ 2.917 \ 0.8264 \ : \ 58.33] \times 0.9143 = [0 \ 2.667 \ 0.7556 \ : \ 53.33]$$

Subtract the result from
Equation 3

$$\begin{array}{r} [0 \ 2.667 \ 0.5556 \ : \ 53.10] \\ - [0 \ 2.667 \ 0.7556 \ : \ 53.33] \\ \hline [0 \ 0 \ -0.2 \ : \ -0.23] \end{array}$$

Substitute new equation for
Equation 3

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.917 & 0.8264 & : 58.33 \\ 0 & 0 & -0.2 & : -0.23 \end{array} \right]$$

Back Substitution

Back Substitution

$$\left[\begin{array}{ccc|c} 144 & 12 & 1 & : 279.2 \\ 0 & 2.917 & 0.8264 & : 58.33 \\ 0 & 0 & -0.2 & : -0.23 \end{array} \right] \Rightarrow \left[\begin{array}{ccc|c} 144 & 12 & 1 & a_1 \\ 0 & 2.917 & 0.8264 & a_2 \\ 0 & 0 & -0.2 & a_3 \end{array} \right] = \left[\begin{array}{c} 279.2 \\ 58.33 \\ -0.23 \end{array} \right]$$

Solving for a_3

$$-0.2a_3 = -0.23$$

$$a_3 = \frac{-0.23}{-0.2} = 1.15$$

Back Substitution (cont.)

$$\begin{bmatrix} 144 & 12 & 1 \\ 0 & 2.917 & 0.8264 \\ 0 & 0 & -0.2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 279.2 \\ 58.33 \\ -0.23 \end{bmatrix}$$

Solving for a_2

$$2.917a_2 + 0.8264a_3 = 58.33$$

$$\begin{aligned} a_2 &= \frac{58.33 - 0.8264a_3}{2.917} \\ &= \frac{58.33 - 0.8264 \times 1.15}{2.917} \\ &= 19.67 \end{aligned}$$

Back Substitution (cont.)

$$\begin{bmatrix} 144 & 12 & 1 \\ 0 & 2.917 & 0.8264 \\ 0 & 0 & -0.2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 279.2 \\ 58.33 \\ -0.23 \end{bmatrix}$$

Solving for a_1

$$144a_1 + 12a_2 + a_3 = 279.2$$

$$\begin{aligned} a_1 &= \frac{279.2 - 12a_2 - a_3}{144} \\ &= \frac{279.2 - 12 \times 19.67 - 1.15}{144} \\ &= 0.2917 \end{aligned}$$

Gaussian Elimination with Partial Pivoting Solution

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 106.8 \\ 177.2 \\ 279.2 \end{bmatrix}$$

$$\begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 0.2917 \\ 19.67 \\ 1.15 \end{bmatrix}$$

Gauss Elimination with Partial Pivoting Another Example

<http://numericalmethods.eng.usf.edu>

Partial Pivoting: Example

Consider the system of equations

$$10x_1 - 7x_2 = 7$$

$$-3x_1 + 2.099x_2 + 6x_3 = 3.901$$

$$5x_1 - x_2 + 5x_3 = 6$$

In matrix form

$$\begin{bmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 3.901 \\ 6 \end{bmatrix}$$

Solve using Gaussian Elimination with Partial Pivoting using five significant digits with chopping

Partial Pivoting: Example

Forward Elimination: Step 1

Examining the values of the first column

$|10|$, $|-3|$, and $|5|$ or 10, 3, and 5

The largest absolute value is 10, which means, to follow the rules of Partial Pivoting, we switch row1 with row1.

Performing Forward Elimination

$$\left[\begin{array}{ccc|c} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right] = \left[\begin{array}{c} 7 \\ 3.901 \\ 6 \end{array} \right] \implies \left[\begin{array}{ccc|c} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 2.5 & 5 \end{array} \right] \left[\begin{array}{c} x_1 \\ x_2 \\ x_3 \end{array} \right] = \left[\begin{array}{c} 7 \\ 6.001 \\ 2.5 \end{array} \right]$$

Partial Pivoting: Example

Forward Elimination: Step 2

Examining the values of the first column

$|-0.001|$ and $|2.5|$ or 0.0001 and 2.5

The largest absolute value is 2.5 , so row 2 is switched with row 3

Performing the row swap

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & -0.001 & 6 \\ 0 & 2.5 & 5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 6.001 \\ 2.5 \end{bmatrix} \implies \begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & -0.001 & 6 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.001 \end{bmatrix}$$

Partial Pivoting: Example

Forward Elimination: Step 2

Performing the Forward Elimination results in:

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & 0 & 6.002 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.002 \end{bmatrix}$$

Partial Pivoting: Example

Back Substitution

Solving the equations through back substitution

$$\begin{bmatrix} 10 & -7 & 0 \\ 0 & 2.5 & 5 \\ 0 & 0 & 6.002 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 7 \\ 2.5 \\ 6.002 \end{bmatrix}$$
$$x_3 = \frac{6.002}{6.002} = 1$$
$$x_2 = \frac{2.5 - 5x_3}{2.5} = -1$$
$$x_1 = \frac{7 + 7x_2 - 0x_3}{10} = 0$$

Partial Pivoting: Example

Compare the calculated and exact solution

The fact that they are equal is coincidence, but it does illustrate the advantage of Partial Pivoting

$$[X]_{calculated} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix} \quad [X]_{exact} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} 0 \\ -1 \\ 1 \end{bmatrix}$$

THE END

<http://numericalmethods.eng.usf.edu>

Determinant of a Square Matrix Using Naïve Gauss Elimination Example

<http://numericalmethods.eng.usf.edu>

Theorem of Determinants

If a multiple of one row of $[A]_{n \times n}$ is added or subtracted to another row of $[A]_{n \times n}$ to result in $[B]_{n \times n}$ then $\det(A) = \det(B)$

Theorem of Determinants

The determinant of an upper triangular matrix

$[A]_{n \times n}$ is given by

$$\det(A) = a_{11} \times a_{22} \times \dots \times a_{ii} \times \dots \times a_{nn}$$

$$= \prod_{i=1}^n a_{ii}$$

Forward Elimination of a Square Matrix

Using forward elimination to transform $[A]_{n \times n}$ to an upper triangular matrix, $[U]_{n \times n}$.

$$[A]_{n \times n} \rightarrow [U]_{n \times n}$$

$$\det(A) = \det(U)$$

Example

Using naïve Gaussian elimination find the determinant of the following square matrix.

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix}$$

Forward Elimination

Forward Elimination: Step 1

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix}$$

Divide Equation 1 by 25 and multiply it by 64, $\frac{64}{25} = 2.56$.

$$[25 \ 5 \ 1] \times 2.56 = [64 \ 12.8 \ 2.56]$$

$$\begin{bmatrix} 64 & 8 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 64 & 12.8 & 2.56 \end{bmatrix}$$

$$\begin{bmatrix} 0 & -4.8 & -1.56 \end{bmatrix}$$

Subtract the result from Equation 2

Substitute new equation for Equation 2

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 144 & 12 & 1 \end{bmatrix}$$

Forward Elimination: Step 1 (cont.)

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 144 & 12 & 1 \end{bmatrix}$$

Divide Equation 1 by 25 and multiply it by 144, $\frac{144}{25} = 5.76$.

$$[25 \ 5 \ 1] \times 5.76 = [144 \ 28.8 \ 5.76]$$

$$\begin{array}{r} [144 \ 12 \ 1] \\ - [144 \ 28.8 \ 5.76] \\ \hline [0 \ -16.8 \ -4.76] \end{array}$$

Subtract the result from Equation 3

Substitute new equation for Equation 3

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & -16.8 & -4.76 \end{bmatrix}$$

Forward Elimination: Step 2

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & -16.8 & -4.76 \end{bmatrix}$$

Divide Equation 2 by -4.8
and multiply it by -16.8 ,
 $\frac{-16.8}{-4.8} = 3.5$.

$$([0 \quad -4.8 \quad -1.56]) \times 3.5 = [0 \quad -16.8 \quad -5.46]$$

Subtract the result from
Equation 3

$$\begin{array}{r} [0 \quad -16.8 \quad -4.76] \\ - [0 \quad -16.8 \quad -5.46] \\ \hline [0 \quad 0 \quad 0.7] \end{array}$$

Substitute new equation for
Equation 3

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix}$$

Finding the Determinant

After forward elimination

$$\left[\begin{array}{ccc} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{array} \right]$$

$$\begin{aligned}\det(A) &= u_{11} \times u_{22} \times u_{33} \\ &= 25 \times (-4.8) \times 0.7 \\ &= -84.00\end{aligned}$$

Summary

- Forward Elimination
- Back Substitution
- Pitfalls
- Improvements
- Partial Pivoting
- Determinant of a Matrix

Additional Resources

For all resources on this topic such as digital audiovisual lectures, primers, textbook chapters, multiple-choice tests, worksheets in MATLAB, MATHEMATICA, MathCad and MAPLE, blogs, related physical problems, please visit

http://numericalmethods.eng.usf.edu/topics/gaussian_elimination.html

THE END

<http://numericalmethods.eng.usf.edu>