

## Chapter 08.04

### Runge-Kutta 4th Order Method for Ordinary Differential Equations-More Examples

#### Chemical Engineering

##### Example 1

The concentration of salt  $x$  in a home made soap maker is given as a function of time by

$$\frac{dx}{dt} = 37.5 - 3.5x$$

At the initial time,  $t = 0$ , the salt concentration in the tank is 50 g/L Using Runge-Kutta 4<sup>th</sup> order method and a step size of,  $h = 1.5$  min, what is the salt concentration after 3 minutes?

##### Solution

$$\frac{dx}{dt} = 37.5 - 3.5x$$

$$f(t, x) = 37.5 - 3.5x$$

$$x_{i+1} = x_i + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)h$$

For  $i = 0$ ,  $t_0 = 0$ ,  $x_0 = 50$

$$\begin{aligned}k_1 &= f(t_0, x_0) \\ &= f(0, 50) \\ &= 37.5 - 3.5(50) \\ &= -137.5\end{aligned}$$

$$\begin{aligned}k_2 &= f\left(t_0 + \frac{1}{2}h, x_0 + \frac{1}{2}k_1h\right) \\ &= f\left(0 + \frac{1}{2}(1.5), 50 + \frac{1}{2}(-137.5)(1.5)\right) \\ &= f(0.75, -53.125) \\ &= 37.5 - 3.5(-53.125) \\ &= 223.44\end{aligned}$$

$$\begin{aligned}k_3 &= f\left(t_0 + \frac{1}{2}h, x_0 + \frac{1}{2}k_2h\right) \\ &= f\left(0 + \frac{1}{2}(1.5), 50 + \frac{1}{2}(223.44)(1.5)\right)\end{aligned}$$

$$\begin{aligned}
&= f(0.75, 217.58) \\
&= 37.5 - 3.5(217.58) \\
&= -724.02 \\
k_4 &= f(t_0 + h, x_0 + k_3 h) \\
&= f(0 + 1.5, 50 + (-724.03)1.5) \\
&= f(1.5, -1036.0) \\
&= 37.5 - 3.5(-1036.0) \\
&= 3663.6 \\
x_1 &= x_0 + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)h \\
&= 50 + \frac{1}{6}(-137.5 + 2(223.44) + 2(-724.02) + (3663.6))1.5 \\
&= 50 + \frac{1}{6}(2525.0)1.5 \\
&= 681.24 \text{ g/L}
\end{aligned}$$

$x_1$  is the approximate concentration of salt at

$$\begin{aligned}
t &= t_1 = t_0 + h = 0 + 1.5 = 1.5 \\
x(1.5) &\approx x_1 = 681.24 \text{ g/L}
\end{aligned}$$

For  $i = 1$ ,  $t_1 = 1.5$ ,  $x_1 = 681.24$

$$\begin{aligned}
k_1 &= f(t_1, x_1) \\
&= f(1.5, 681.24) \\
&= 37.5 - 3.5(681.24) \\
&= -2346.8
\end{aligned}$$

$$\begin{aligned}
k_2 &= f\left(t_1 + \frac{1}{2}h, x_1 + \frac{1}{2}k_1 h\right) \\
&= f\left(1.5 + \frac{1}{2}1.5, 681.24 + \frac{1}{2}(-2346.8)1.5\right) \\
&= f(2.25, -1078.9) \\
&= 37.5 - 3.5(-1078.9) \\
&= 3813.6
\end{aligned}$$

$$\begin{aligned}
k_3 &= f\left(t_1 + \frac{1}{2}h, x_1 + \frac{1}{2}k_2 h\right) \\
&= f\left(1.5 + \frac{1}{2}1.5, 681.24 + \frac{1}{2}(3813.6)1.5\right) \\
&= f(2.25, 3541.4) \\
&= 37.5 - 3.5(3541.4) \\
&= -12358
\end{aligned}$$

$$k_4 = f(t_1 + h, x_1 + k_3 h)$$

$$\begin{aligned}
&= f(1.5 + 1.5, 681.24 + (-12358)1.5) \\
&= f(3, -17855) \\
&= 37.5 - 3.5(-17855) \\
&= 62530 \\
x_2 &= x_1 + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)h \\
&= 681.24 + \frac{1}{6}(-2346.8 + 2(3813.6) + 2(-12358) + 62530)1.5 \\
&= 681.24 + \frac{1}{6}(43096)1.5 \\
&= 11455 \text{ g/L}
\end{aligned}$$

$x_2$  is the approximate concentration of salt at

$$\begin{aligned}
t_2 &= t_1 + h = 1.5 + 1.5 = 3 \text{ min} \\
x(3) &\approx x_2 = 11455 \text{ g/L}
\end{aligned}$$


The exact solution of the ordinary differential equation is given by

$$x(t) = 10.714 + 39.286e^{-3.5t}$$

The solution to this nonlinear equation at  $t = 3$  min is

$$x(3) = 10.715 \text{ g/L}$$

Figure 1 compares the exact solution with the numerical solution using Runge-Kutta 4<sup>th</sup> order method using different step sizes.




**Figure 1** Comparison of Runge-Kutta 4<sup>th</sup> order method with exact solution for different step sizes.

Table 1 and Figure 2 show the effect of step size on the value of the calculated temperature at  $t = 3$  min.


**Table 1** Value of concentration of salt at 3 minutes for different step sizes.

Step size, $h$	$x(3)$	$E_t$	$ \epsilon_t  \%$
3	14120	-14109	131680
1.5	11455	-11444	106800
0.75	25.559	-14.843	138.53
0.375	10.717	-0.0014969	0.013969
0.1875	10.715	-0.00031657	0.0029544



**Figure 2** Effect of step size in Runge-Kutta 4<sup>th</sup> order method.

In Figure 3, we are comparing the exact results with Euler's method (Runge-Kutta 1<sup>st</sup> order method), Heun's method (Runge-Kutta 2<sup>nd</sup> order method) and Runge-Kutta 4<sup>th</sup> order method.



**Figure 3** Comparison of Runge-Kutta methods of 1<sup>st</sup>, 2<sup>nd</sup>, and 4<sup>th</sup> order.